

Mersenne Binomials and the Coefficients of the non-associative Exponential

by L. Gerritzen

Abstract

The non-associative exponential series $\exp(x)$ is a power series with monomials from the magma M of finite, planar rooted trees. The coefficient $a(t)$ of $\exp(x)$ relative to a tree t of degree n is a rational number and it is shown that

$$\hat{a}(t) := \frac{a(t)}{2^{n-1} \cdot \prod_{i=1}^{n-1} (2^i - 1)}$$

is an integer which is a product of Mersenne binomials. One obtains summation formulas

$$\sum \hat{a}(t) = \omega(n)$$

where the sum is extended over all trees t in M of degree n and

$$\omega(n) = \frac{2^{n-1}}{n!} \prod_{i=1}^{n-1} (2^i - 1).$$

The prime factorization of $\omega(n)$ is described. The sequence $(\omega(n))_{n \geq 1}$ seems to be of interest.

Introduction

For a natural number n , we denote by M_n the n -th Mersenne number $2^n - 1$.

The Mersenne factorial $n!_M$ is defined to be the product $\prod_{i=1}^n M_i$ of all Mersenne number $M_1, \dots, M_n$ while the Mersenne binomial $\binom{n}{r}_M$ is defined to be

$$\frac{n!_M}{r!_M (n-r)!_M}$$

for natural numbers r between 0 and n .

In quantum calculus, see for instance [KC], there are the notions $[n]$ for the q - analogue of $n \in \mathbb{N}$, $[n]!$ for the q - factorial and $\begin{bmatrix} n \\ r \end{bmatrix}$ for the q - binomial coefficients which are polynomials in the variable q . If $f(2)$ denotes the value of a polynomial f after substituting 2 for q , then $[n](2) = M_n$, $[n]!(2) = n!_M$ and $\begin{bmatrix} n \\ r \end{bmatrix}(2) = \binom{n}{r}_M$.

Let M be the magma with unit 1_M freely generated by a single element x . It can be identified with the set of all finite planar binary rooted trees, see [DG].

Let $\mathbb{Q}\{\{x\}\}$ the $\mathbb{Q}$ - algebra of power series with monomials from M . It was proved in [DG] that there is a unique series $\exp(x) \in \mathbb{Q}\{\{x\}\}$ such that $\exp(x) = 1 + x + \text{higher terms}$ and

$$\exp(x) \cdot \exp(x) = \exp(2x)$$

see also Proposition (3.2). Let $a(t)$ be the coefficient of the non-associative exponential $\exp$ relative to the planar binary rooted tree t and put

$$\hat{a}(t) = n! \cdot \omega(n) \cdot a(t)$$

where

$$\omega(n) := \frac{2^{n-1}(n-1)!_M}{n!}.$$

A main result of this article states that all $\hat{a}(t)$ are integers which are obtained as products of Mersenne binomials. The projection of the non-associative exponential to the classical one leads to the decomposition

$$\sum \hat{a}(t) = \omega(n)$$

if the sum is extended over all trees of degree n .

In section 1 the notions of Mersenne order and Wieferich exponents at odd primes is defined. One gets results about the prime factorization of Mersenne numbers.

In section 2 we show that all $\omega(n)$ are integers by computing the order $\text{ord}_p \omega(n)$ for all primes p .

The applications about the coefficients $a(t)$ of $\exp(x)$ are deduced in section 3.

I would like to thank Doron Zeilberger for helpful hints.

1 Mersenne orders

We call $M_n = 2^n - 1$ the n -th Mersenne number for $n \in \mathbb{N}, n \geq 1$.

This is in contrast to the definition by D. Shanks, see [S], Chap. 1, which is also widely used in which M_n is called Mersenne number only if n is a prime.

Let p be an odd prime.

Definition 1.1. The Mersenne order $v(p)$ at p is the smallest number $r \in \mathbb{N}, r \geq 1$, such that p divides M_r .

As $2^{v(p)} \equiv 1 \pmod{p}$ and $v(p)$ is the order of the class of 2 in $(\mathbb{Z}/p\mathbb{Z})^*$, we get that $v(p)$ divides the order of the multiplicative group $(\mathbb{Z}/p\mathbb{Z})^*$ which is $(p-1)$.

Also p divides M_n , if and only if n is a multiple of $v(p)$.

Also $p < 2^{v(p)}$ from which follows that $\log_2(p) < v(p)$.

It has the consequence that $\lim_{p \rightarrow \infty} v(p) = \infty$.

Definition 1.2. Let $\varepsilon(p) := \text{ord}_p(M_{v(p)})$ for any odd prime; it is called the Wieferich exponent at p .

Always $\varepsilon(p) \geq 1$ and p is called a Wieferich prime if $\varepsilon(p) \geq 2$.

Up to now only two Wieferich primes are known, namely 1093 and 3511 and it was checked that these are the only Wieferich primes $< 4 \cdot 10^{12}$ in [CDP].

The Mersenne order at 1093 is $364 = \frac{1092}{3} = 2^2 \cdot 7 \cdot 13$ and the Mersenne order at 3511 is $1755 = \frac{3510}{2} = 3^3 \cdot 5 \cdot 13$.

The Wieferich exponent for both primes is equal to 2.

Proposition 1.3. Let p be an odd prime, $n \in \mathbb{N}, n \geq 1$.

- (i) $\text{ord}_p(M_n) = 0$, if n is not a multiple of $v(p)$.
- (ii) If $n = v(p) \cdot p^r \cdot k$ and p does not divide k , the $\text{ord}_p(n) = r$ and $\text{ord}_p(M_n) = \varepsilon(p) + r$

Proof. Statement (i) is well-known.

Let now $n = v(p) \cdot p^r \cdot k$ as in (ii).

Let m be an integer and consider the group $(\mathbb{Z}/p^m\mathbb{Z})^*$ of units of the ring $\mathbb{Z}/p^m\mathbb{Z}$.

Let U_m be the subgroup of $(\mathbb{Z}/p^m\mathbb{Z})^*$ of class congruent to 1 modulo p .

Then U_m has order p^{m-1} because any element α in U_m has a unique representation

by a number $a = 1 + \sum_{i=1}^{m-1} a_i p^i$ with $a_i \in \{0, \dots, p-1\}$. The order of α is $m - \min(\{i : a_i \neq 0\}, m)$ because if $a = 1 + p^\varepsilon \cdot x$, x any number not divisible by p , then

$$a^p = 1 + p^{\varepsilon+1} \cdot y$$

where

$$y = x + \sum_{i=2}^p \binom{p}{i} x^i p^{i\varepsilon - \varepsilon - 1}$$

which shows that $y \equiv x \pmod{p}$ as $p > 2$.

Let now $\alpha = \bar{2}^{v(p)}$ where $\bar{2}$ is the class of 2 in $\mathbb{Z}/p^m\mathbb{Z}$.

Then $\alpha \in U_m$ and α is represented by a number $a = 1 + p^{\varepsilon(p)} \cdot x$, where x is not divisible by p . Then α^{p^r} is represented by a number

$$1 + p^{\varepsilon(p)+r} \cdot y$$

with y not divisible by p .

It follows that $\text{ord}_p M_n = \varepsilon(p) + r$ because the order of α^{p^r} is equal to the order of α^{p^k} . $\square$

Corollary 1.4.

$$M_n = \prod_n p^{\varepsilon(p) + \text{ord}_p(n)}$$

where the product is extended over all odd primes p for which the Mersenne order $v(p)$ divides n .

2 Factorial Mersenne quotients

$F(n) := \prod_{i=1}^n M_i$ is called the Mersenne factorial of $n \in \mathbb{N}$; it is also denoted by $n!_M$.

Let

$$\omega(n) := \frac{2^{n-1} \cdot F(n-1)}{n!}$$

for $n \geq 1$; it is called the factorial Mersenne quotient at n . Let $m \in \mathbb{N}, m \geq 1$, and p be a prime. There is a unique p -adic expansion

$$m = \sum_{i=0}^r m_i p^i$$

with $m_i \in \{0, 1, \dots, p-1\}$. Let

$$d_p(m) := \sum_{i=0}^r m_i.$$

Proposition 2.1. (i) $\text{ord}_2 \omega(n) = d_2(n) - 1$

(ii) If p is an odd prime, then

$$\text{ord}_p(\omega(n)) = \varepsilon(p) \cdot m - \frac{(n - d_p(n)) - (m - d_p(m))}{(p-1)}$$

where $v(p)$ is the Mersenne order at p , m is the largest integer $\leq (n-1)/v(p)$ and $\varepsilon(p)$ is the Wieferich exponent at p .

Proof. 1) $m \equiv d_p(m) \pmod{p-1}$ because $m \equiv \sum_{i=0}^r m_i \pmod{p-1}$. It is well known that

$$\text{ord}_p(n!) = \frac{n - d_p(n)}{(p-1)} = \sum_{i=1}^{<\infty} \left[\frac{n}{p^i} \right]$$

1) $\text{ord}_2(\omega(n)) = (n-1) - \text{ord}_2(n!)$ because $F(n)$ is odd. Thus

$$\text{ord}_2(\omega(n)) = d_2(n) - 1 \geq 0$$

for $n \geq 1$.

2) Let now p be an odd prime. Then

$$\text{ord}_p F(n-1) = \sum_{i=1}^{n-1} \text{ord}_p M_i.$$

If $i = k \cdot p^r v(p)$ and k is not divisible by p , then

$$\text{ord}_p M_i = \varepsilon(p) + r$$

and $\text{ord}_p M_i = 0$ if p is not divisible by p .

$$\text{ord}_p(F(n-1)) = \sum_{j=1}^m \text{ord}_p M_{jv(p)}$$

where $m_p(n) = m := \left[\frac{n-1}{v(p)} \right]$ is the largest integer $\leq (n-1)/v(p)$. It follows from Proposition 1.3 that

$$\text{ord}_p M_{j\sigma(p)} = \varepsilon(p) + \text{ord}_p(j)$$

as $\text{ord}_p(j) = \text{ord}_p(jv(p))$. As

$$\sum_{j=1}^m \text{ord}_p(j) = \text{ord}_p(m!) = \frac{m - d_p(m)}{p-1}$$

we get the formulas

$$\text{ord}_p F(n-1) = \varepsilon(p) \cdot m + \frac{m - d_p(m)}{p-1}$$

and

$$\text{ord}_p(\omega(n)) = \varepsilon(p)m - \frac{(n - d_p(n)) - (m - d_p(m))}{p-1}$$

□

Proposition 2.2. $\omega(n) \in \mathbb{N}$.

Proof. We will show now that $\text{ord}_p \omega(n) \geq 0$. From the formula in 3) we get

$$\begin{aligned} \text{ord}_p(\omega(n)) &\geq m - \frac{(n - d_p(n)) - (m - d_p(m))}{p - 1} \\ &\geq m - \frac{n - d_p(n)}{(p - 1)} \geq \left[\frac{n - 1}{p - 1} \right] - \frac{n - d_p(n)}{p - 1} \geq 0 \end{aligned}$$

because $m - d_p(m) \geq 0$, $m \geq \frac{n-1}{p-1}$, $n - d_p(n) \leq n - 1$ and $\frac{n-d_p(n)}{p-1} \in \mathbb{N}$.

(3) From 1) and 2) we get that $\text{ord}_p(\omega(n - 1)) \geq 0$ for all primes p . As $\omega(n - 1)$ is a rational number, it follows that $\omega(n - 1)$ is an integer. $\square$

Example 2.3. (1) $\text{ord}_7 \omega(13) = 3$, because $n = 13, m = \frac{13-1}{v(7)} = 4, d_7(4) = 4, d_7(13) = 7, \varepsilon(7) = 1, n - d_7(n) = 6, m - d_7(m) = 0$ $\text{ord}_7(\omega(13)) = 4 - \frac{6}{6} = 3$
(2) $\text{ord}_7 \omega(100) = 21$, because $v(7) = 3, n = 100, m = \left[\frac{n-1}{3} \right] = 33, d_7 = 100) = 4, d_7(33) = 9$ $\text{ord}_7(\omega(100)) = 33 - \frac{96-24}{6} = 21$, because $\log(\omega(n)) \approx \binom{n+1}{2} \log(2) + n - (n + 1/2) \log(n)$.

Let $\pi_M(x)$ be defined to be number of odd primes p such that the Mersenne order $v(p)$ is $\leq x - 1$.

It seems interesting to determine the asymptotic behaviour of π_M .

One can expect some results about π_M from the prime factorization of $n!_M$, because $\pi_M(n)$ is the number of primes dividing $n!_M$.

Example 2.4. $\pi_M(16) = 15$ because the primes dividing $16!_M$ are the odd primes ≤ 16 and the primes 17, 23, 31, 43, 73, 89, 127, 151, 257, 8191.

3 Coefficients of the non-associative exponential

Let M be the magma with neutral element 1_M freely generated by a set consisting of one element x . The multiplication in M is a map $\cdot : M \times M \rightarrow M$ and the restriction of this map onto $(M - \{1_M\}) \times (Mx\{1_M\})$ is injective and its image is $M - \{1_M, x\}$. It means that for any $t \in M - \{1_M, x\}$ there is a unique pair $(t_1, t_2) \in M \times M, t_1 \neq 1_M \neq t_2$, such that $t = t_1 \cdot t_2$.

There is a unique homomorphism $\deg : M \rightarrow \mathbb{N}$ such that $\deg(1_M) = 0, \deg(x) = 1$.

It is important to realize that any $t \in M, t \neq 1_M$, gives rise to a unique planar binary rooted tree with $\deg(t)$ leaves, see [1], section 1, p. 163. The grafting of such trees corresponds to the multiplication map of M .

Let $P = \mathbb{Q}\{\{x\}\}$ be the $\mathbb{Q}$ -algebra of power series with monomials from M . Thus any $f \in P$ has a unique expansion

$$f = \sum_{t \in M} c(t) \cdot t$$

with $c(t) \in \mathbb{Q}$. It will also be called the algebra of tree power series in x or the algebra of power series over $\mathbb{Q}$ in a nonassociative and noncommutative variable x .

Define the order $\text{ord}(f)$ of f to be $\min\{\deg(t) : c(t) \neq 0\}$, if $f \neq 0$ and $\text{ord}(0) = +\infty$. The following proposition is obtained by standard calculus methods.

Proposition 3.1. (i) *There is a unique $\mathbb{Q}$ -linear map $\frac{d}{dx} : P \rightarrow P$ such that:*

$$\frac{d}{dx}(x) = 1$$

and

$$\frac{d}{dx}(f \cdot g) = \frac{d}{dx}(f) \cdot g + f \cdot \frac{d}{dx}(g)$$

for all $f, g \in P$.

$f' := \frac{d}{dx}(f)$ is called the derivative of f with respect to x .

(ii) *Let $g \in P$ with $\text{ord}(g) \geq 1$. Then there is a unique $\mathbb{Q}$ -algebra homomorphism $\eta_g : P \rightarrow P$ such that $\eta_g(x) = g$. One denotes $\eta_g(x)$ also by $f \circ g$ or $f(g)$ and calls η_g the substitution homomorphism induced by g .*

(iii) *There is a canonical $\mathbb{Q}$ -algebra homomorphism $[\] : P \rightarrow \mathbb{Q}[[x]]$, where $\mathbb{Q}[[x]]$ denotes the classical $\mathbb{Q}$ -algebra of power series in an associative (and commutative) variable x such that $[x] = x$.*

One calls $[f]$ the classical power series associated to $f \in P$.

Proposition 3.2. *There is a unique $f \in \mathbb{Q}\{\{x\}\}$ such that*

$$f'(0) = 1$$

$$f(x) \cdot f(x) = f(2x)$$

Moreover $f'(x) = f(x)$ where $f'(x)$ denotes the derivative of f relative to x . The tree power series $f(x)$ is called the nonassociative, noncommutative exponential series and will be denoted in this article by e^x or $\exp(x)$.

Proof. Inductively we define a map $a : M \rightarrow \mathbb{Q}$ by putting $a(t) := 1$, if $t = 1_M$ or $t = x$ and otherwise

$$a(t) := \frac{a(t_1) \cdot a(t_2)}{2^n - 2}$$

if $n = \deg(t)$ and $t = t_1 \cdot t_2$ with $t_i \in M$.

Put $f := \sum_{t \in M} a(t) \cdot t \in P = \mathbb{Q}\{\{x\}\}$. It is easy to check that $f(x) \cdot f(x) = f(2x)$:

$$\begin{aligned} f(2x) &= \sum_{t \in M} a(t) \cdot 2^{\deg(t)} \cdot t \\ f(x) \cdot f(x) &= \sum_{(t_1, t_2) \in M \times M} a(t_1) a(t_2) t_1 \cdot t_2 \\ &= 1 + 2a(x) + \\ &\quad + \sum_{t \in M, \deg(t) \geq 2, t = (t_1, t_2)} (a(t) \cdot a(x^0) + a(x^0) a(t) + a(t_1) a(t_2)) \cdot t. \end{aligned}$$

As $2a(t) + a(t_1) \cdot a(t_2) = 2^n a(t)$ for any $t \in M$ with $\deg(t) \geq 2, t = t_1 \cdot t_2$, we obtain the desired functional equation.

For the proof that $f'(x) = f(x)$ we refer to [DG]. $\square$

From the proof we get that the coefficients $a(t)$ of $\exp$ satisfy: $a(1) = a(x) = 1$ and

$$a(t_1 \cdot t_2) = \frac{a(t_1) a(t_2)}{2^n - 2}$$

where $n = \deg(t_1 \cdot t_2)$.

Let $\hat{a}(t) := n \cdot \omega(n) \cdot a(t)$.

Proposition 3.3. *Let $t_1, t_2 \in M, n_i = \deg(t_i) \geq 1$ and $t = t_1 \cdot t_2, n = \deg(t) = n_1 + n_2$. Then*

$$\hat{a}(t) = \left(\frac{n-2}{n_1-1}\right)_M \hat{a}(t_1) \hat{a}(t_2)$$

Proof.

$$\begin{aligned} \hat{a}(t) &= n! \cdot \omega(n) \cdot \frac{a(t_1) a(t_2)}{2^n - 2} \\ &= n! \cdot \frac{\omega(n)}{2M_{n-1}} \frac{\hat{a}(t_1)}{n_1! \omega(n_1)} \frac{\hat{a}(t_2)}{n_2! \omega(n_2)} \\ &= \frac{2^{n-1} F(n-1)}{2M_{n-1}} \frac{\hat{a}(t_1) \hat{a}(t_2)}{2^{n_1-1} F(n_1-1) 2^{n_2-1} F(n_2-1)} \\ &= \frac{F(n-2)}{F(n_1-1) F(n_2-1)} \hat{a}(t_1) \hat{a}(t_2) \\ &= \left(\frac{n-2}{n_1-1}\right)_M \hat{a}(t_1) \hat{a}(t_2). \end{aligned}$$

$\square$

It follows from (3.3) that $\hat{a}(t)$ is a product of Mersenne binomials. More precisely:

Corollary 3.4. $\hat{a}(t) = \prod_{a \in I(t)} \binom{n(a) - 2}{n_1(a) - 1}_M \in \mathbb{N}$ where $I(t)$ is the set of inner nodes of t and $n(a)$ is the degree of $t_{\leq a}$ where $t_{\leq a}$ is the tree below a which is defined to consist of all nodes b for which the simple path from b to the root of t is passing through a . Also $n_1(a)$ is the degree of the left factor s_1 of $t_{\leq a}$ (such that $t_{\leq a} = s_1 \cdot s_2$).

Example 3.5. Let T_n be the set of trees in M of degree n defined as follows: $T_1 = \{x\}$.

If T_{n-1} is already defined, then $T_n = x \cdot T_{n-1} \cup T_{n-1} \cdot x$.

Then $\#T_n = 2^{n-2}$ for $n \geq 2$. One can show that $\hat{a}(t) = 1$ iff $t \in T_n$.

Corollary 3.6.

$$\sum_{\deg(t)=n} \hat{a}(t) = \omega(n)$$

Proof. We consider the canonical algebra homomorphism

$$[\] : \mathbb{Q}\{\{x\}\} \rightarrow \mathbb{Q}[[x]].$$

Then $[exp]$ is the classical series $\sum_{n=0}^{\infty} \frac{x^n}{n!}$. As $[t] = x^n$ if $t \in M$ has degree n , it

follows that $\sum_{\deg(t)=n} \hat{a}(t) = \frac{1}{n!}$. □

For $1 \leq k \leq n-1$ let

$$S_k(n) = \sum_{\substack{\deg(t_1)=k \\ \deg(t_2)=n-k}} \hat{a}(t_1 \cdot t_2).$$

Then

$$S_k(n) = \binom{n-2}{k-1}_M \sum_{\substack{\deg(t_1)=k \\ \deg(t_2)=n-k}} \hat{a}(t_1) \hat{a}(t_2)$$

$$S_k(n) = \binom{n-2}{k-1}_M \cdot \omega(k) \cdot \omega(n-k)$$

It follows from Corollary 3.6 that $\omega(n) = \sum_{k=1}^{n-1} S_k(n)$. Thus

Corollary 3.7. $\sum_{k=1}^{n-1} \binom{n-2}{k-1}_M \omega(k) \cdot \omega(n-k) = \omega(n)$.

Example 3.8. $\omega(1) = \omega(2) = 1, \omega(3) = 2, \omega(4) = 7, \omega(5) = 42 = 2 \cdot 3 \cdot 7$
Thus

$$\begin{aligned}\omega(6) &= \sum_{k=1}^5 \binom{4}{k-1}_M \omega(k) \omega(6-k) \\ &= 1 \cdot \omega(5) + M_4 \cdot 1 \cdot \omega(4) + \frac{M_4 \cdot M_3}{M_2} \omega(3)^2 + M_4 \cdot \omega(4) \cdot 1 + 1 \cdot \omega(5) \\ &= 2 \cdot 42 + 30 \cdot 7 + 5 \cdot 7 \cdot 4 = 434 = 2 \cdot 7 \cdot 31\end{aligned}$$

How to understand from this partition of $\omega(6)$ that 31 is a factor of $\omega(6)$?

References

- [CDP] *Crandall, R. - Dilcher, K. - Pomerance, C.* : A search for Wieferich and Wilson primes, Math. Comp. 66 (1997), no. 217, p. 433 - 449
- [DG] *Drensky, V.- Gerritzen, L.*: On non-associative exponential and logarithm, (2002), to appear in Journal of Algebra
- [KC] *Kac, V. - Cheung, P.*: Quantum Calculus, Springer-Verlag, Universitext, 2002
- [S] *Shanks, D.*: Solved and unsolved problems in number theory, Third edition, Chelsea Publishing Co., New York 1985

Key Words

Binary rooted trees, non-associative power series, exponential series, q -binomials, Mersenne binomials

Address: Prof. Dr. Lothar Gerritzen

Ruhr-Universität Bochum

Fakultä für Mathematik

D 44780 Bochum

Germany

Tel.-number: 0049-0234-32-28304

FAX: 0049-0234-32-14025

Email: Lothar.Gerritzen@rub.de